

## Dosage

Visit [Motrin.com](https://www.motrin.com) for additional product information

mL = milliliter

### Infants' MOTRIN® Concentrated Drops

### Children's MOTRIN® Oral Suspension

### Children's MOTRIN® Chewables

### MOTRIN® IB Caplets

### MOTRIN® IB Liquid Gels

### MOTRIN® IB Migraine Liquid Gels

*Chew or crush tablets completely before swallowing*

Active ingredient:  
ibuprofen (NSAID)\*  
50 mg (in each 1.25 mL)

\*Nonsteroidal anti-inflammatory drug.

Active ingredient:  
ibuprofen (NSAID)\*  
100 mg (in each 5 mL)

\*Nonsteroidal anti-inflammatory drug.

Active ingredient:  
ibuprofen (NSAID)\*  
100 mg (in each chewable tablet)

\*Nonsteroidal anti-inflammatory drug.

Active ingredient:  
ibuprofen (NSAID)\*  
200 mg (in each caplet)

\*Nonsteroidal anti-inflammatory drug.

Active ingredient:  
ibuprofen (NSAID)\*  
200 mg (in each capsule)

\*Nonsteroidal anti-inflammatory drug.

Active ingredient:  
ibuprofen (NSAID)\*  
200 mg (in each capsule)

\*Nonsteroidal anti-inflammatory drug.

Weight (lbs)

Age

12-17 lbs      6-11 mos

1.25 mL

Infants' MOTRIN® and Children's MOTRIN® are different concentrations. The markings on the included dosing cup correspond to the OTC label doses listed below.

-

18-23 lbs      12-23 mos

1.875 mL

-

24-35 lbs      2-3 years

5 mL

1 tablet

36-47 lbs      4-5 years

7.5 mL

1½ tablets

48-59 lbs      6-8 yrs

The OTC label does not include dosing directions for children ages 2 years and older.

10 mL

2 tablets

60-71 lbs      9-10 yrs

12.5 mL

2½ tablets

72-95 lbs      11 yrs

15 mL

3 tablets

OTC label does not include dosing for children under 12 years old.

OTC label does not include dosing for children under 12 years old.

OTC label does not include dosing for children under 18 years old.

If needed, repeat dose every 6-8 hours  
Do **NOT** use more than 4 times a day

If needed, repeat dose every 6-8 hours  
Do **NOT** use more than 4 times a day

If needed, repeat dose every 6-8 hours  
Do **NOT** use more than 4 times a day

#### OTC LABEL STATES:

**Adults and children 12 years and older:**  
Take 1 caplet every **4 to 6 hours** while symptoms persist; if pain or fever does not respond to 1 caplet, 2 caplets may be used

**Children under 12 years:**  
Ask a doctor

Do **NOT** take more than 6 caplets in **24 hours** unless directed by a doctor

#### OTC LABEL STATES:

**Adults and children 12 years and older:**  
Take 1 capsule every **4 to 6 hours** while symptoms persist; if pain or fever does not respond to 1 capsule, 2 capsules may be used

**Children under 12 years**  
Ask a doctor

Do **NOT** take more than 6 capsules in **24 hours** unless directed by a doctor

#### OTC LABEL STATES:

**Adults:**  
Take 2 capsules with a glass of water; if symptoms persist or worsen, ask your healthcare professional

**Under 18 years of age:**  
Ask a doctor  
Do **NOT** take more than 2 capsules in **24 hours** unless directed by a doctor

#### Remind parents & caregivers:

Read & follow the label on all MOTRIN® products.

If needed, repeat dose every 6-8 hours while symptoms last.

Do **NOT** give more than 4 doses in 24 hours.

Use only the dosing device (syringe or dosing cup) that came with the product.

\*Nonsteroidal anti-inflammatory drug

Use product only as directed.

© Johnson & Johnson Consumer Inc. 2023

#### Important Instructions For Proper Use:

Always read and follow the label on all MOTRIN® products. Keep carton.

Wait the right amount of time between doses

If you have any questions, ask your healthcare professional which product may be right for you

\*Nonsteroidal anti-inflammatory drug